



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

III. Solution by G. W. GREENWOOD, M. A. (Oxon), Lebanon, Ill.

$$\Pi(a\beta - \gamma\delta) = a\beta\gamma\delta(\Sigma a)^2 - (\Sigma a\beta\gamma)^2 = m^2(-a)^2 - (-am)^2 = 0.$$

Hence the product of any one pair of the roots equals that of the remaining pair.

Solved by A. H. Holmes by actual computation of the roots.

Also solved by the Proposer.

207. Proposed by A. J. PAULSEN, San Francisco, Cal.

$$\text{Solve } x^4 + y^4 = 2x^2y^2; \quad x + y = a.$$

Solution by EDWIN L. RICH, Lehigh University, and A. H. HOLMES, Brunswick, Maine.

Writing the first of these equations in the form $(x+y)^2(x-y)^2 = 0$, it is easily seen, since $x+y=a$, that the four values of x are ∞ , $-\infty$, $\frac{1}{2}a$, $\frac{1}{2}a$; and those of y are $-\infty$, ∞ , $\frac{1}{2}a$, $\frac{1}{2}a$.

Also solved by G. B. M. Zerr, G. W. Greenwood, M. E. Graber, and J. Scheffer.

GEOMETRY.

233. Proposed by S. F. NORRIS, Professor of Mathematics, Baltimore City College, Baltimore, Md.

If from any point on a circle circumscribed about a triangle perpendiculars are dropped to the sides of the triangle, the feet of these perpendiculars lie on a line. [Ashton's *Plane and Solid Analytic Geometry*, page 87, 11th example].

Remark by G. W. GREENWOOD, M. A. (Oxon), Lebanon, Ill.

See solution of Geometry Problem number 184, August-September, 1902.

234. Proposed by M. E. GRABER, A. B., Instructor in Mathematics and Physics in Heidelberg University, Tiffin, Ohio.

Find the curve which is reciprocal to a circle and define it as a locus.

I. Solution by the PROPOSER.

If A be the center of the given circle, P any point on it; x and a lines corresponding to P and A ; and X the point ax , $AP = \frac{OA \cdot OX}{Ox} \sin ax$. Denoting OX by z the ratio $\sin ax : \sin xz$ is constant. The reciprocal to a circle is then the envelope of a line x which divides the angle between a fixed line a and a variable line z passing through a fixed point O , into parts whose sines are in a constant ratio. Defined as a locus, the reciprocal curve to a circle is the path of a point which moves so that its distance from a fixed point varies as its distance from a fixed straight line. [Lachlan's *Modern Pure Geometry*.]

II. Solution by G. W. GREENWOOD, M. A. (Oxon), Professor of Mathematics and Astronomy in McKendree College, Lebanon, Ill.

See Russell's *Pure Geometry*, Chapter VIII, §11, or Salmon's *Conic Sections*, §308. The following is an analytic solution. Call the center of the given circle C and its radius a . Call the center of the circle of reciprocation O . De-